

# 2026 Zorvec™ active application advice for potatoes.



Products containing Zorvec™ active deliver highly effective disease control against potato Late Blight.

Corteva strongly advise following the 2026 application advice when recommending and applying Zorvec products.



## Best practice application advice for 2026



1. Apply Zorvec Entecta in strict alternation with effective chemistry of a different mode of action, other than a CAA (benthiavalcarb, mandipropamid, dimethomorph).
2. The post-Zorvec application interval to the following non-Zorvec containing spray should not exceed 7 days. 
3. In curative situations (disease evident or confirmed in local area) Zorvec co-formulations should be applied in tank-mix with a partner fungicide containing a curative active ingredient e.g. cymoxanil, (Option®) or propamocarb.
4. Sprays containing Zorvec should not exceed 20% of the total number of sprays applied to a crop. 
5. A single crop should not receive more than 2 sprays containing Zorvec products.
6. Only 1 application would be applied to a seed crop as part of the late blight programme.

## Resources to help you

- Visit our Zorvec active hub to go back to basics on why this chemistry and its application advice is so important: [www.corteva.com/ie/zorvec](http://www.corteva.com/ie/zorvec)
- Discover Corteva's range of products to help you maximise your potato yield at: [www.corteva.com/ie/crops/potatoes](http://www.corteva.com/ie/crops/potatoes)

## Need further advice?

Speak to your advisor or our Technical Services team who are always at hand to help answer any questions you may have.

**Hotline:**

**+44 800 689 8899**

**Email:**

**[ukhotline@corteva.com](mailto:ukhotline@corteva.com)**

Product containing Zorvec active:

**Zorvec Entecta™**  
FUNGICIDE

# Zorvec™ active