(Container)

Enlist Duo™
Herbicide

with COLEX-D™ technology

GROUP	4		9	HERBICIDES

For control of annual and perennial weeds and use in Enlist™ field corn, Enlist™ E3 soybeans, summerfallow, and prior to seeding or after seeding (but before crop emergence) in spring and winter varieties of wheat (including durum), barley, rye and field corn

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: 2,4-D, present as choline salt 194 g /L
glyphosate, present as dimethylamine salt 204 g /L
Solution

REGISTRATION NO. 30958 PEST CONTROL PRODUCTS ACT

CAUTION – EYE IRRITANT
POTENTIAL SKIN SENSITIZER

NET CONTENTS: 1 L - Bulk

Corteva Agriscience Canada Company
Suite 240, 115 Quarry Park Rd. SE
Calgary, AB
T2C 5G9
1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS
KEEP OUT OF REACH OF CHILDREN
CAUSES IRRITATION OF SKIN AND EYES
DO NOT GET IN EYES. CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY
POTENTIAL SKIN SENSITIZER
AVOID CONTACT WITH SKIN AND CLOTHING
DO NOT APPLY BY AIR

DO NOT eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, or using the toilet. Vehicle cabs must be decontaminated regularly.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

Restricted Entry Interval (REI)

DO NOT enter or allow worker entry into treated areas during the intervals specified in the following table:

Crop	Postapplication Activity	Restricted Entry Interval (REI)
All crops	Hand line irrigation	6 days
	Scouting	24 hours
	All other activities	12 hours

PROTECTIVE CLOTHING AND EQUIPMENT

Follow the personal protective equipment, engineering controls and restriction requirements for the appropriate mixer/loader and applicator scenario as described below. These restrictions are required to minimize exposure to the worker. Application may need to be performed over multiple days or using multiple applicators.

Application Equipment and Engineering Controls	Personal Protective Equipment		Restricted Amount of Product Handled per Day per Person
	Mixer/Loader / Clean-up and Repair	Applicator	
Groundboom – Open Cab	Coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eyewear (goggles or face shield).	Coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.	Less than 1803 L
	Chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eyewear (goggles or face shield).	Chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.	No restriction

Groundboom – Closed Cab[1]	Coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eyewear (goggles or face shield).	Long-sleeved shirt, long pants, socks and shoes.	No restriction
Mechanically-pressurized handgun - fine to very coarse spray[2]	Chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eyewear (goggles or face shield).		Less than 34 L
	Chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes, protective eyewear (goggles or face shield) and respirator[3].		Less than 45 L
All other handheld application equipment - fine to very coarse spray[2]	Coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eyewear (goggles or face shield).		No restriction

[1] Gloves are not required within a closed cab.
[2] Based on the American Society of Agricultural and Biological Engineers (ASABE) ASABE S572.1 Droplet Size Classification.
[3] A respirator is a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides or a NIOSH-approved canister approved for pesticides.

If the product comes in contact with clothing/PPE, remove all contaminated clothing/PPE, wash with soap and water and dress in clean clothing. Wash the outside of gloves with soap and water before removing. Launder contaminated clothing/PPE regularly and separately from other laundry.

Refer to Directions for Use section for any additional requirements.

MIXING
Transfer System
For containers larger than 20 L, use a transfer system that avoids open pouring when transferring the liquid concentrate from such containers into the spray tank.

PHYSICAL OR CHEMICAL HAZARDS
Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fibreglass, plastic and plastic-lined steel containers. DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

FIRST AID
Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.
If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.
If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION
No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient. This product may cause severe irritation to the eyes. Overexposure to 2,4-D may cause coughing, burning, dizziness or temporary loss of muscle coordination. Other possible effects of overexposure include fatigue, muscle weakness or nausea. Treat symptomatically.

AGRICULTURAL CHEMICAL
DO NOT ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS
Toxic to birds and small wild mammals
Toxic to aquatic organisms and non-target terrestrial plants.
Observe spray buffer zones specified under DIRECTIONS FOR USE.
This product will harm other broadleaved plants in the vicinity of the treatment area. If applying this product using a handheld sprayer, DO NOT directly spray or allow the spray to drift onto ornamentals or gardens.
DO NOT spray exposed roots of trees and ornamentals.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated area into aquatic habitats avoid application to areas with moderate to steep slope, compacted soil, or clay. Avoid application of this product when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website

STORAGE
Store this product away from food or feed. This product must be stored away from fertilizers, seeds, insecticides, fungicides or other herbicides intended for use on 2,4-D and glyphosate sensitive crops.

DISPOSAL
Recyclable Containers
DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

If this product is exposed to temperatures below 0°C, it should be warmed to at least 5°C and mixed thoroughly before using.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

(Booklet)

Enlist Duo™ Herbicide

with COLEX-D™ technology

GROUP	4	9	HERBICIDES

For control of annual and perennial weeds and use in Enlist™ field corn, Enlist™ E3 soybeans, summerfallow, and prior to seeding or after seeding (but before crop emergence) in spring and winter varieties of wheat (including durum), barley, rye and field corn

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT:	2,4-D, present as choline salt	194 g /L
	glyphosate, present as dimethylamine salt	204 g /L

Solution

REGISTRATION NO. 30958 PEST CONTROL PRODUCTS ACT

CAUTION – EYE IRRITANT
POTENTIAL SKIN SENSITIZER

NET CONTENTS: 1 L - Bulk

Corteva Agriscience Canada Company
Suite 240, 115 Quarry Park Rd. SE
Calgary, AB
T2C 5G9
1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS
KEEP OUT OF REACH OF CHILDREN
CAUSES IRRITATION OF SKIN AND EYES
DO NOT GET IN EYES. CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY
POTENTIAL SKIN SENSITIZER
AVOID CONTACT WITH SKIN AND CLOTHING
DO NOT APPLY BY AIR

DO NOT eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, or using the toilet. Vehicle cabs must be decontaminated regularly.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

Restricted Entry Interval (REI)

DO NOT enter or allow worker entry into treated areas during the intervals specified in the following table:

Crop	Postapplication Activity	Restricted Entry Interval (REI)
All crops	Hand line irrigation	6 days
	Scouting	24 hours
	All other activities	12 hours

PROTECTIVE CLOTHING AND EQUIPMENT

Follow the personal protective equipment, engineering controls and restriction requirements for the appropriate mixer/loader and applicator scenario as described below. These restrictions are required to minimize exposure to the worker. Application may need to be performed over multiple days or using multiple applicators.

Application Equipment and Engineering Controls	Personal Protective Equipment		Restricted Amount of Product Handled per Day per Person
	Mixer/Loader / Clean-up and Repair	Applicator	
Groundboom – Open Cab	Coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eyewear (goggles or face shield).	Coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.	Less than 1803 L
	Chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eyewear (goggles or face shield).	Chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.	No restriction

Groundboom – Closed Cab[1]	Coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eyewear (goggles or face shield).	Long-sleeved shirt, long pants, socks and shoes.	No restriction
Mechanically-pressurized handgun - fine to very coarse spray[2]	Chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eyewear (goggles or face shield).		Less than 34 L
	Chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes, protective eyewear (goggles or face shield) and respirator[3].		Less than 45 L
All other handheld application equipment - fine to very coarse spray[2]	Coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eyewear (goggles or face shield).		No restriction

[1] Gloves are not required within a closed cab.
[2] Based on the American Society of Agricultural and Biological Engineers (ASABE) ASABE S572.1 Droplet Size Classification.
[3] A respirator is a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides or a NIOSH-approved canister approved for pesticides.

If the product comes in contact with clothing/PPE, remove all contaminated clothing/PPE, wash with soap and water and dress in clean clothing. Wash the outside of gloves with soap and water before removing. Launder contaminated clothing/PPE regularly and separately from other laundry.

Refer to Directions for Use section for any additional requirements.

MIXING
Transfer System
For containers larger than 20 L, use a transfer system that avoids open pouring when transferring the liquid concentrate from such containers into the spray tank.

PHYSICAL OR CHEMICAL HAZARDS
Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fibreglass, plastic and plastic-lined steel containers. DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

FIRST AID
Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.
If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION
No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient. This product may cause severe irritation to the eyes. Overexposure to 2,4-D may cause coughing, burning, dizziness or temporary loss of muscle coordination. Other possible effects of overexposure include fatigue, muscle weakness or nausea. Treat symptomatically.

AGRICULTURAL CHEMICAL
DO NOT ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS
Toxic to birds and small wild mammals
Toxic to aquatic organisms and non-target terrestrial plants.
Observe spray buffer zones specified under DIRECTIONS FOR USE.
This product will harm other broadleaved plants in the vicinity of the treatment area. If applying this product using a handheld sprayer, do not directly spray or allow the spray to drift onto ornamentals or gardens.
DO NOT spray exposed roots of trees and ornamentals.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with moderate to steep slope, compacted soil, or clay.
Avoid application of this product when heavy rain is forecast.
Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website

STORAGE
Store this product away from food or feed. This product must be stored away from fertilizers, seeds, insecticides, fungicides or other herbicides intended for use on 2,4-D and glyphosate sensitive crops.

DISPOSAL

Recyclable Containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

If this product is exposed to temperatures below 0°C, it should be warmed to at least 5°C and mixed thoroughly before using.

DIRECTIONS FOR USE

USE PRECAUTIONS

- As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
- DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application

DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) coarse to extremely coarse classification. Boom height must be 60 cm or less above the crop or ground.

Glyphosate is not to be applied using hand-wicking or hand-daubing methods.

Ground Application - Cropland

Apply 50 to 200 L of spray solution per hectare depending on the type of application equipment used. Use sufficient water for even distribution. Spray at low pressures (200 to 275 kPa) when the weeds are actively growing.

Droplet Size: Apply as a coarse to extremely coarse spray (ASABE S572 (572.1 to 572.3) Standard). Use drift reducing nozzle tips in accordance with manufacturer directions that produce a droplet classification of coarse to extremely coarse to significantly reduce the potential for drift.

Refer to the Product Use Guide for further information.

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of Enlist Duo Herbicide.
4. Fill the sprayer tank with sufficient water (see water volumes in Ground Application - Cropland section). Maintain sufficient agitation in the spray tank during mixing and spraying to ensure a uniform spray mixture.
5. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops.
6. Follow sprayer clean-up directions as listed (see Sprayer Clean-Out Instructions section).

Spot Treatment

For backpack application of Enlist Duo Herbicide for spot treatment of weeds such as thistles, mix 0.2 L of product in 10 L of water. Wet all foliage thoroughly. Maximum 2 applications per year.

Maximum Amount of Product Applied/day/person with Handheld Equipment: 14.5 L.

Tank Mixtures

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Corteva Agriscience Canada Company at 1-800-667-3852 or www.corteva.ca for information before applying any tank mix that is not specifically recommended on this label.

Drift Control Additives

Enlist Duo Herbicide contains drift control technology. If desired, use only a Corteva Agriscience-approved drift control additive. When a drift control additive is used, read and carefully observe the precautionary statements and all other information appearing on the additive label.

FOR USE IN ENLIST™ FIELD CORN ONLY

Enlist Duo Herbicide is a systemic herbicide and is intended for control of emerged annual and perennial weeds. Enlist Duo Herbicide is selective to Enlist field corn, application of Enlist Duo Herbicide to non-Enlist field corn will cause serious crop damage and yield loss.

Rates of application

Use the recommended rate depending on the weeds present (see weed list below).
For harder-to-control weeds, such as common milkweed and yellow nutsedge, Enlist Duo may be tank mixed with a product containing 480 g/L glyphosate, present as the dimethylamine salt (e.g., VP480

Herbicide) at 2.52 or 1.92 L/ha, respectively. Refer to the glyphosate product label for additional information, including spray volumes and weeds controlled at the 3.75 L/ha (1800 g a.e./ha) rate.

The tank mix with glyphosate may be applied a maximum of once per season and only where there is no application of Enlist Duo alone in the same year.

Timing of Application
Apply up to V8 growth stage or 120 cm (48") in height, whichever occurs first. Make 1 to 2 applications with a minimum of 12 days between applications. A second application may be used for late weed flushes emerging after the initial application but must be made before the V8 growth stage. Where there is a mix of susceptible, less susceptible, and/or hard-to-control weeds, use the appropriate rate depending on weed spectrum.

2.9 L/ha will control the following susceptible weeds:

barley, volunteer	lamb's-quarters
bluebur	mustards (except dog and green tansy)
burdock (before 4 leaf)	pigweed, Russian
canola, volunteer♦	pigweed, redroot
chickweed, common	plantain, common
cocklebur	ragweed, common
false flax	ragweed, giant
fleabane, Canada	Russian thistle
fleabane, daisy	shepherd's purse
flixweed	stinkweed
foxtail (giant, green)	sunflower, annual
goatsbeard	sweet clover
hempnettle	vetch
hoary cress	wheat, volunteer
horsetail, field	wild radish
kochia	wild oats

♦including glyphosate tolerant and Clearfield canola varieties.

4.3 L/ha will control the above listed weeds, plus the following less susceptible weeds:

barnyard grass	milkweed, common♦♦
biennial wormwood♦	mustard, dog
bindweed, field♦♦	narrow-leaved hawk's beard
bindweed, hedge	nightflowering catchfly
blue lettuce♦	nightshade, Eastern black flowering
buckwheat, tartary	nutsedge, yellow♦♦
buckwheat, wild	oak leaf goosefoot
burdock♦	palmer amaranth♦♦
Canada thistle♦♦	pigweed, smooth
corn spurry	pineappleweed
cleavers, common	proso millet, wild
cow cockle	purslane, common
crabgrass, smooth	quackgrass
crabgrass, large	round-leaved mallow♦♦
dandelion	smartweed, green
chickweed, mouse-eared♦	smartweed, Pennsylvania
fall panicum	sow-thistle, annual
field peppergrass	sow-thistle, perennial♦♦
hairy galinsoga	tansy, common
knotweed (before 4 leaf)	velvetleaf
lady's thumb	waterhemp, common
leafy spurge♦	wild tomato
glyphosate-resistant common waterhemp♦♦♦	glyphosate-resistant Canada fleabane♦♦♦

♦Top growth control only.

♦♦Use 2 applications for best control. The 2nd application should be no later than the V8 stage of corn. Milkweed should be 15-60 cm in height and actively growing. Yellow nutsedge should be 5-15 cm in height and actively growing. Canada thistle and perennial sow-thistle should be from the rosette stage to 50 cm in height and actively growing. The 2nd application should be at least 2 weeks after the first for best control of these perennial weeds.

♦♦♦For heavy populations, or populations at a more advanced growth stage use the higher rate listed below.

5.5 L/ha will control the above listed weeds, plus the following hard-to-control weeds:

palmer amaranth♦	

♦Hard-to-control weeds may require a total program approach including a soil-applied residual herbicide(s) followed by a single or sequential post herbicide application.

Pre-Harvest Interval

- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.

Restrictions

- Do not apply more than two post emergent applications per use season.
- Do not apply more than 8.6 L/ha of Enlist Duo Herbicide (3.44 kg acid equivalent) per use season.
- Do not apply more than the total seasonal rate of 1800 g a.i./ha glyphosate
- Read and follow the Enlist Stewardship Program (https://www.enlist.com/ca-en/experience-enlist/stewardship.html) that accompanies the use of Enlist field corn seed.

FOR USE IN ENLIST™ E3 SOYBEAN ONLY

Enlist Duo Herbicide is a systemic herbicide and is intended for control of emerged annual and perennial weeds. Enlist Duo Herbicide is selective to Enlist E3 soybeans. Application of Enlist Duo Herbicide to non-Enlist E3 soybeans will cause serious crop damage and yield loss.

Rates of application
Use the recommended rate depending on the weeds present (see weed list below).

For harder to control weeds, such as milkweed, yellow nutsedge and field bindweed, Enlist Duo may be tank mixed with a product containing 480 g/L glyphosate, present as the dimethylamine salt (e.g., VP480 Herbicide) at 2.52 or 1.92 L/ha, respectively. Refer to the glyphosate product label for additional information, including spray volumes and weeds controlled at the 3.75 L/ha (1800 g a.e./ha rate).

The tank mix with glyphosate may be applied a maximum of once per season and only where there is no application of Enlist Duo alone in the same year.

Timing of Application
Apply when weeds are small and soybean growth stage is no later than R2 (full flowering stage). Make one to two applications with a minimum of 12 days between applications. Do not apply Enlist Duo Herbicide to soybeans later than the R2 stage. Where there is a mix of susceptible, less susceptible, and hard-to-control weeds, use the appropriate rate depending on weed spectrum.

2.9 L/ha will control the following susceptible weeds:

barley, volunteer	lamb's-quarters
bluebur	mustards (except dog and green tansy)
burdock (before 4 leaf)	pigweed, Russian
canola, volunteer♦	pigweed, redroot
chickweed, common	plantain, common
cocklebur	ragweed, common
false flax	ragweed, giant
fleabane, Canada	Russian thistle
fleabane, daisy	shepherd's purse
flixweed	stinkweed
foxtail (giant, green)	sunflower, annual
goatsbeard	sweet clover
hempnettle	vetch
hoary cress	wheat, volunteer
horsetail, field	wild radish
kochia	wild oats

♦including glyphosate tolerant and Clearfield canola varieties.

4.3 L/ha will control the above listed weeds, plus the following less susceptible weeds:

barnyard grass	milkweed, common♦♦
biennial wormwood♦	mustard, dog
bindweed, field♦♦	narrow-leaved hawk's beard
bindweed, hedge	nightflowering catchfly
blue lettuce♦	nightshade, Eastern black flowering
buckwheat, tartary	nutsedge, yellow♦♦
buckwheat, wild	oak leaf goosefoot
burdock♦	palmer amaranth♦♦
Canada thistle♦♦	pigweed, smooth
corn spurry	pineappleweed
cleavers, common	proso millet, wild
cow cockle	purslane, common
crabgrass, smooth	quackgrass
crabgrass, large	round-leaved mallow♦♦
dandelion	smartweed, green
chickweed, mouse-eared♦	smartweed, Pennsylvania
fall panicum	sow-thistle, annual
field peppergrass	sow-thistle, perennial♦♦
hairy galinsoga	tansy, common
knotweed (before 4 leaf)	velvetleaf
lady's thumb	waterhemp, common
leafy spurge♦	wild tomato
glyphosate-resistant common waterhemp♦♦♦	glyphosate-resistant Canada fleabane♦♦♦

♦Top growth control only.

♦♦Use 2 applications for best control. The 2nd application should be no later than the V8 stage of corn. Milkweed should be 15-60 cm in height and actively growing. Yellow nutsedge should be 5-15 cm in height and actively growing. Canada thistle and perennial sow-thistle should be from the rosette stage to 50 cm in height and actively growing. The 2nd application should be at least 2 weeks after the first for best control of these perennial weeds.

♦♦♦For heavy populations, or populations at a more advanced growth stage use the higher rate listed below.

5.5 L/ha will control the above listed weeds, plus the following hard-to-control weeds:

palmer amaranth♦	

♦Hard-to-control weeds may require a total program approach including a soil-applied residual herbicide(s) followed by a single or sequential post herbicide application.

Control of volunteer Enlist Corn in Enlist E3 Soybean

A tank mix of Enlist Duo Herbicide at 4.3 L/ha plus either Select Emulsifiable Concentrate Postemergence Herbicide or Centurion Emulsifiable Concentrate Post-Emergence Herbicide at 0.19 L/ha or Poast Ultra Liquid Emulsifiable Herbicide at 0.32 L/ha can be used to control volunteer Enlist Corn in Enlist E3 Soybean.

Pre-Harvest Interval

- Do not graze treated soybeans.
- Do not harvest for forage or hay.

Restrictions

- Do not apply more than two post emergent applications per use season.
- Do not apply more than 8.6 L/ha of Enlist Duo Herbicide (3.44 kg acid equivalent) per use season.
- Do not apply more than the total seasonal rate of 1800 g a.i./ha glyphosate

- Read and follow the Enlist Stewardship Program (https://www.enlist.com/ca-en/experience-enlist/stewardship.html) that accompanies the use of Enlist E3 soybean seed.

FOR USE PRIOR TO SEEDING OR AFTER SEEDING (BUT BEFORE CROP EMERGENCE) IN SPRING AND WINTER VARIETIES OF WHEAT (INCLUDING DURUM), BARLEY, RYE AND FIELD CORN

Enlist Duo Herbicide may be applied to control both annual and perennial weed species prior to seeding or after seeding **(BUT BEFORE CROP EMERGENCE)** in spring and winter varieties of wheat (including durum), barley, rye and field corn.

Apply at a rate of 2.2 L per hectare in a water volume of 50 to 100 L of clean water per hectare as a broadcast spray for control of weeds (up to 8 cm in height) or at a rate of 3.3 L per hectare for control of weeds (up to 15 cm in height). An application rate of 4.3 L/ha in the fall will provide control of red clover.

2.2 L/ha will control the following weeds (up to 8 cm):

bluebur	narrow-leaved hawk's beard
brome, downy	nettle, stinging
burdock	Persian darnel
canola, volunteer♦	pigweed, redroot
cereals, volunteer	pigweed, Russian
cocklebur	plantain, common
flax, false	prickly lettuce
flax, volunteer	radish, wild
fleabane, Canada	ragweed, common
fleabane, daisy	ragweed, false
flixweed	ragweed, giant
foxtail, giant	shepherd's purse
foxtail, green	stinkweed
goat's beard	sunflower, wild
hempnettle	sweet clover
kochia	thistle, Russian
lady's-thumb	thyme-leaved spurge
lamb’s quarters	wild buckwheat♦♦♦
mustards♦♦	wild oats
mustard, wild	

♦1- to 4-leaf stage only, all biotypes
♦♦except dog and tansy mustard
♦♦♦1- to 2-leaf stage only

3.3 L/ha will control the above listed weeds, plus the following weeds (up to 15 cm):

canola, volunteer♦	peppergrass
chickweed, common	pigweed, prostrate
galinsoga, hairy	pigweed, tumble
goosefoot, oak-leaved	pineapple weed
groundsel, common	purslane, common
hawkweed	smartweed
heal-all	sorrel, sheep
knotweed	sowthistle, annual
mustard, dog	velvetleaf
mustard, tansy	wild buckwheat♦♦

♦1- to 6-leaf stage, all biotypes
♦♦1- to 4-leaf stage

4.3 L/ha will control the following weeds:

red clover

FOR USE IN SUMMERFALLOW

Apply at a rate of 3.3 L per hectare in a water volume of 50 to 100 L of clean water per hectare as a broadcast spray for control of weeds (up to 15 cm in height).

3.3 L/ha will control the following weeds (up to 15 cm)

bluebur	narrow-leaved hawk's beard
brome, downy	nettle, stinging
burdock	peppergrass
canola, volunteer♦	Persian darnel
cereals, volunteer	pigweed, prostrate
chickweed, common	pigweed, redroot
cocklebur	pigweed, Russian
flax, false	pigweed, tumble
flax, volunteer	pineapple weed
fleabane, Canada	plantain, common
fleabane, daisy	prickly lettuce
flixweed	purslane, common
foxtail, giant	radish, wild
foxtail, green	ragweed, common
galinsoga, hairy	ragweed, false
goat's beard	ragweed, giant
goosefoot, oak-leaved	shepherd's purse
groundsel, common	smartweed
hawkweed	sorrel, sheep
heal-all	sowthistle, annual
hempnettle	stinkweed
knotweed	sunflower, wild

kochia	sweet clover
lady's-thumb	thistle, Russian
lamb's quarters	thyme-leaved spurge
mustard, dog	velvetleaf
mustard, tansy	wild buckwheat♦♦
mustard, wild	wild oats
mustards	

♦1- to 6-leaf stage, all biotypes
♦♦1- to 4-leaf stage

Sprayer Clean-Out Instructions
To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

1. Immediately after spraying, completely drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. First rinse:
   - Spray the inside of tank with clean water and fill the sprayer with at least one tenth of the spray tank volume.
   - Agitate and circulate for 15 minutes, and flush through booms and hoses.
   - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
   - Drain tank completely.
3. Second rinse:
   - Fill the tank with clean water.
   - Add All Clear Spray Tank Decontaminator, or Clean-Out Spray Tank Cleaner, or 1 L of household ammonia (containing a minimum of 3 % ammonia) per 100 L of water, or similar tank cleaning agent as per manufacturer's recommendations while filling the tank with clean water.
   - Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full. Allow to stand for 15 minutes with agitation. Flush the solution out of the spray tank through the spray booms. Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
   - If possible, let the solution stand in the sprayer tank and booms for an extended period of time, overnight if possible.
   - After flushing the boom and hoses, drain tank completely.
   - Remove nozzles and screens and clean separately with a cleaning agent or an ammonia solution (100 mL in 10 L water).
4. Third rinse:
   - Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume.
   - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
   - Drain tank completely.

Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty odour which may cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

Sensitive Plants
Vegetables, flowers, grapes, fruit trees and other desirable plants are sensitive to 2,4-D and glyphosate, even in minute quantities. Care should be taken to avoid spraying these types of plants or allowing spray mist to drift onto these plants during both their growing and dormant periods. Coarse sprays are less likely to drift. At higher temperatures, vaporization may cause injury to susceptible plants growing nearby.

This product may cause damage to lawns or pastures if applied before the grass is well established. In addition, most legumes may be damaged or killed.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,
- low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, wetlands and estuarine/marine water bodies.

Method of Application	Crop	Maximum number of applications	Spray Buffer Zones (metres) Required for the Protection of:	
			Aquatic Habitats	Terrestrial Habitats
Field sprayer	Enlist field corn	2	1	2
	Enlist E3 soybeans	4	1	2
	Summerfallow, spring and winter wheat (including durum), barley, rye and field corn (pre-seeding treatments only)	1	1	1
	Spring and winter wheat (including durum), barley, rye and field corn (after seeding)	2	1	2

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that Enlist Duo Herbicide is both a Group 4 and a Group 9 herbicide. Any weed population may contain or develop plants naturally resistant to Group 4 and/or Group 9 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Enlist Duo Herbicide or other Group 4 and/ or Group 9 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.

- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Corteva Agriscience Canada Company at 1-800-667-3852 or at www.corteva.ca.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

™Trademark of Corteva Agriscience and its affiliated companies.

011526

Label Code: CN-30958-011-E
Replaces: CN-30958-010-E

Specimen Label Notes
Added control of GR Canada fleabane, GR waterhemp and palmer amaranth. Increased rate 5.5L/ha