OcTTain™ XL

HERBICIDE

Trusted, easy to use, high performance broadleaf weed control.

WHY USE OCTTAIN™ XL HERBICIDE?

- **Performance and value.** The standard for kochia control (including Group 2 and Group 9 resistant), plus wild buckwheat and 25 other key broadleaf weeds.
- **Ease of use.** Available in 40-acre cases, 240-acre drums and 1,280-acre totes.
- **Crop-safe and flexible.** Excellent crop safety in barley, wheat and durum.
- Tank mix with Simplicity™ GoDRI™ (wheat), Liquid Achieve™ (barley) herbicides, and other grass weed control products.

BROADLEAF WEEDS CONTROLLED

- Annual sunflower
- Blue lettuce[1, 4]
- Bluebur
- Burdock
- Cleavers[3]
- Cocklebur
- Dandelion[2, 4]
- Docks[4]
- Dog mustard[4]
- Field bindweed[1, 4]
- Field horsetail[1]
- Field peppergrass[4]
- Flixweed
- Goat's-beard
- Gumweed[4]
- Hairy galinsoga[4]
- Hedge bindweed[4]
- Hemp nettle
- Hoary cress[1]
- Kochia[3]
- Lamb's-quarters
- Leafy spurge[1, 4]
- Mustards *(except green and grey tansy)*
- Oak-leaved goosefoot[4]
- Plantain
- Prickly lettuce
- Ragweed
- Round-leaved mallow
- Russian thistle[4]
- Shepherd's purse
- Smartweed[4]
- Stinkweed
- Stork's-bill
- Sweet clover
- Tansy mustard[4]
- Tartary buckwheat[4]
- Vetch
- Volunteer canola[5]
- Volunteer flax
- Wild buckwheat
- Wild radish

WEEDS SUPPRESSED

- Annual sow thistle[4]
- Canada thistle[1, 4]
- Chickweed[3]
- Perennial sow thistle[1]

1 Top growth control only.
2 Spring rosettes.
3 Including Group 2 resistant biotypes.
4 Requires the addition of 86 mL/ac (2 oz/ac) of 2,4-D Ester 700.
5 All herbicide-tolerant varieties.

APPLICATION GUIDELINES

CROPS	Barley Durum wheat Spring wheat Winter wheat
PACKAGING	**Case:** 2 x 9 L jugs **Pallet:** 5 x 108 L drums **Tote:** 576 L
RATES	450 mL/ac
ACRES TREATED	• 20 ac/jug • 240 ac/drum • 1280 ac/tote
WATER VOLUME	• Ground 20-40 L/ac (5-10 US gal/ac) • Aerial 12-20 L/ac (3-5 US gal/ac)
TIMING	**Crop stage:** 4-leaf to just prior to flag-leaf emergence **Weed stage:** 1 to 6-leaf
RAINFAST	1 hour
TANK-MIX HERBICIDES	• Everest® • Horizon® • Liquid Achieve™ SC herbicide • Puma® Super • Simplicity™ herbicide • Simplicity™ GoDRI™ herbicide • Traxos®

DOWNLOAD THE 2025 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK-MIX INSTRUCTIONS.

CROP ROTATION

- Alfalfa
- Barley
- Canola
- Corn
- Dry beans
- Faba beans
- Flax
- Forage grasses
- Lentils
- Mustard
- Oats
- Peas
- Potatoes
- Rye
- Soybeans
- Sugar beets
- Sunflowers
- Wheat

GRAZING AND HARVEST

- Allow 7 days after application before grazing lactating animals.
- Withdraw meat animals from treated areas at least 3 days before slaughter.
- Allow 30 days after application before cutting hay or harvesting forage.
- Mature crops may be harvested 60 days after application.

AVAILABLE IN BULK