

HERBICIDE
POST-EMERGENCE

GROUP

4

APPLICATION GUIDELINES

Prominex™

Arylex™ active

HERBICIDE

Prominex™ herbicide with Arylex™ active delivers exceptional control of annual AND perennial broadleaf weeds with the convenience of an all-in-one formulation.

WHY USE PROMINEX™ HERBICIDE?

- **Unmatched Weed Control.** Prominex provides control of both annual AND perennial broadleaf weeds
- **Just GO.** Thanks to Arylex active, you can Just GO on small or large broadleaf weeds, in early or late crop staging, and even in cool or dry conditions
- **Tank Mix Flexibility.** Choose whether to tank mix with a Group 1 grass herbicide in your wheat or barley and a Group 2 grass herbicide in your wheat
- **Convenient Formulation.** All-in-one ME formulation allows for ultimate convenience and easy loading
- **Expanded Group 4 Mode of Action Control.** Tank mix with MCPA Ester 600 or 2,4-D Ester for expanded Group 4 broadleaf weed control

Prominex + 236 mL/ac MCPA Ester 600

GRASS WEEDS CONTROLLED

- Barnyard grass

ANNUAL WEEDS CONTROLLED

- American dragonhead
- Ball mustard
- Burdock
- Canada fleabane**
- Canada thistle
- Chickweed**
- Cleavers**
- Cocklebur
- Common ragweed**
- Cow cockle
- False ragweed
- Flixweed*
- Giant ragweed**
- Hemp-nettle*
- Henbit
- Kochia**
- Lamb's-quarters
- Nightshade species (Eastern black, hairy and cutleaf)
- Plantain
- Prickly lettuce
- Redroot pigweed
- Round-leaved mallow
- Shepherd's purse*
- Stinkweed
- Stork's-bill
- Velvetleaf
- Vetch
- Volunteer alfalfa
- Volunteer canola (all herbicide tolerant varieties)
- Volunteer flax
- Wild buckwheat
- Wild mustard
- Wild radish

WEEDS SUPPRESSED

- Dandelion
- Field horsetail
- Perennial sow-thistle
- Smartweed*



CROPS

- Barley
- Durum wheat
- Spring wheat
- Winter wheat

RATES AND ACRES TREATED

- Rates:**
- 414 mL/ac
- Acres treated:**
- 40 ac/case (20 ac/jug)
- Water volume:**
- 10-20 US gal/ac

PACKAGING

- Case:**
- 2 x 8.3 L jugs

WHEN TO APPLY

- 3-leaf to just prior to flag leaf emergence

RAINFAST

- 4 hours

HERBICIDE TANK MIXES

- MCPA Ester 600 • 2, 4-D Ester 700
- Simplicity™ GoDRI™

Crop rotation

10 months:

- Barley
- Canola
- Corn
- Flax
- Forage grasses
- Oats
- Mustard

22 months:

- Peas¹
- Fall rye (not underseeded with legumes, clover or alfalfa)
- Soybeans¹
- Summerfallow
- Wheat
- Lentils
- Chickpeas

¹For pea or soybean rotation, rainfall from June 1 to August 31 in the year of application must be greater than 140 mm (5.5 inches) and annual rainfall must be greater than 175 mm (6.9 inches).

*Including Group 2 resistant biotypes

**Including Groups 2 and 9 resistant biotypes

For a complete weed list and specific weed staging, please refer to the product label.